arXiv:1510.07826v3 [hep-th] 27 Jan 2016

Scattering of instantons, monopoles and vortices in higher dimensions

Tatiana A. Ivanova

Bogoliubov Laboratory of Theoretical Physics, JINR
141980 Dubna, Moscow Region, Russia

Email: ita@theor.jinr.ru

Abstract

We consider Yang-Mills theory on manifolds $\mathbb{R}\times X$ with a d-dimensional Riemannian manifold X of special holonomy admitting gauge instanton equations. Instantons are considered as particle-like solutions in $d+1$ dimensions whose static configurations are concentrated on X. We study how they evolve in time when considered as solutions of the Yang-Mills equations on $\mathbb{R}\times X$ with moduli depending on time $t\in\mathbb{R}$. It is shown that in the adiabatic limit, when the metric in the X direction is scaled down, the classical dynamics of slowly moving instantons corresponds to a geodesic motion in the moduli space $\mathcal{M}$ of gauge instantons on X. Similar results about geodesic motion in the moduli space of monopoles and vortices in higher dimensions are briefly discussed.

1 Introduction and summary

Instantons in four dimensions are nonperturbative Bogomolny-Prasad-Sommerfield (BPS) configurations solving first-order anti-self-duality equations for gauge fields which imply the full Yang-Mills equations [1]. If one makes the assumption that the non-abelian gauge potential is independent of one coordinate on $\mathbb{R}^4$ then the anti-self-duality equations are reduced to Bogomolny equations on $\mathbb{R}^3$ describing non-abelian monopoles [2]. Furthermore, considering anti-self-dual Yang-Mills equations on a four-manifold $\Sigma_2 \times S^2$ and imposing SO(3)-equivariance condition on gauge fields, one obtains vortex equations on a two-dimensional Riemannian manifold Σ_2 (see e.g. [2, 3] and references therein). Vortices, monopoles and instantons are important objects in modern field theories describing nonperturbative physics [4]-[7].

Non-abelian monopoles are also particle-like static solutions of Yang-Mills-Higgs equations in Minkowski space $\mathbb{R}^{3,1}$ [1]. Vortices can also be obtained as static solutions of Yang-Mills-Higgs equations in $2+1$ dimensions [4]-[7]. Similarly instantons can be considered as solitons in $4+1$ dimensional Yang-Mills theory. One can ask about the dynamics of all these solitons which can evolve according to the second-order field equations of Yang-Mills-Higgs or Yang-Mills theory. In the seminal paper [8] Manton suggested that in the "slow motion limit" the monopole dynamics can be described in terms of geodesics in the moduli space of static multi-monopole solutions. These geodesics are defined via a metric on the multi-monopole moduli space [8]. This heuristic approach was extended to vortices in 2+1 dimensions [9], domain walls in 3+1 dimensions [10] and instantons in $4+1$ dimensions (see e.g. [11]). Higher derivative corrections to the lowest-order (adiabatic) results were considered in [12]. The Manton approach was rigorously justified both for monopoles and vortices by Stuart [13]. However, we are not aware about such a justification for scattering instantons in 4+1 dimensions. Here we provide a derivation of motion of instantons along geodesics in the multi-instanton moduli space by using the adiabatic approach. Generalizing [11], we will consider this approach for instantons not only in four but also in higher dimensions.

Instanton equations on a d-dimensional Riemannian manifold X can be introduced as follows [14]-[16]. Suppose there exists a 4-form Q on X. Then there exists a $(d-4)$-form $\Sigma := *Q$, where $*$ is the Hodge operator on X defined with the help of a metric g on X. Let $\mathcal{A}$ be a connection on a rank-k vector bundle E over X with the curvature $\mathcal{F}_X = \mathrm{d}\mathcal{A} + \mathcal{A}\wedge\mathcal{A}$. For simplicity we choose SU(k) as our gauge group and therefore both $\mathcal{A}$ and $\mathcal{F}_X$ take values in the Lie algebra su(k). The generalized anti-self-duality (instanton) equation on the gauge field $\mathcal{F}_X$ is [16]

$$*\mathcal{F}_X + \Sigma\wedge\mathcal{F}_X \;=\; 0\;. \tag{1.1}$$

For $d > 4$ these equations can be defined on manifolds X with *special holonomy*, i.e. such that the holonomy group G^h of the Levi-Civita connection on the tangent bundle TX is a subgroup in SO(d). Solutions of (1.1) satisfy the Yang-Mills equations

$$D_\mu\mathcal{F}_X^{\mu\nu} := \frac{1}{\sqrt{\det g}}\,\partial_\mu(\sqrt{\det g}\,\mathcal{F}_X^{\mu\nu}) + [\mathcal{A}_\mu, \mathcal{F}_X^{\mu\nu}] \;=\; 0\;, \tag{1.2}$$

where the derivatives $\partial_\mu := \partial/\partial x^\mu$ are taken with respect to local coordinates x^μ on X and $g = g_{\mu\nu}\,\mathrm{d}x^\mu\mathrm{d}x^\nu, \mu, \nu, ... = 1, ..., d$. The instanton equations are also well defined on manifolds X with non-integrable G^h-*structures*, i.e. when $\mathrm{d}\Sigma \neq 0$. In this case (1.1) imply the Yang-Mills equations with (3-form) torsion $T := *\mathrm{d}\Sigma$, as is discussed e.g. in [17]-[20]. Such torsionful Yang-Mills equations naturally appear in heterotic string compactifications with H-flux.

We extend the manifold X by the time axis $\mathbb{R}$ and introduce on the Lorentzian manifold $M = \mathbb{R} \times X$ a metric

$$\hat{g}_\varepsilon = -\mathrm{d}t^2 + \varepsilon^2 g \ , \tag{1.3}$$

where $t = x^0$ is a coordinate on $\mathbb{R}$ and ε is a real parameter. Denoting by $\{x^{\hat{\mu}}\} = \{x^0, x^\mu\}$ local coordinates on $M = \mathbb{R} \times X$, we introduce the Yang-Mills equations on M,

$$\hat{D}_{\hat{\mu}} \hat{\mathcal{F}}^{\hat{\mu}\hat{\nu}} := \frac{1}{\sqrt{\det g}} \, \partial_{\hat{\mu}} (\sqrt{\det g} \, \hat{\mathcal{F}}^{\hat{\mu}\hat{\nu}}) + [\hat{\mathcal{A}}_{\hat{\mu}}, \hat{\mathcal{F}}^{\hat{\mu}\hat{\nu}}] \ = \ 0 \ , \tag{1.4}$$

where we used the fact that $|\det \hat{g}_\varepsilon| = \varepsilon^{2d} \det g$.

It is not easy to construct non-trivial time-dependent solutions of the Yang-Mills equations (1.4). The *adiabatic limit* method, based on Manton's idea, provides a useful and powerful tool for describing such solutions. The adiabatic limit refers to the geometric process of shrinking the metric (1.3) in the X direction by taking the limit $\varepsilon \to 0$. We will show that solutions of the Yang-Mills equations (1.4) in the limit $\varepsilon \to 0$ for the metric (1.3) converge to the solutions of one-dimensional sigma-model describing a map from $\mathbb{R}$ into the moduli space of gauge instantons on X. For connections $\mathcal{A}$ not depending on one coordinate of X we will get geodesics in the moduli space of (generalized) monopoles on a $(d-1)$-dimensional submanifold of X. Similar reductions to geodesic in moduli space of (generalized) vortices on $(d-2)$-dimensional submanifolds of X will also be described.

2 Moduli space of instantons in $d \geq 4$

Moduli space of connections.[1] Let X be an oriented smooth manifold of dimension d, G a semisimple compact Lie group, $\mathfrak{g}$ its Lie algebra, P a principal G-bundle over X, $\mathcal{A}$ a connection 1-form on P and $\mathcal{F}_X = \mathrm{d}\mathcal{A} + \mathcal{A} \wedge \mathcal{A}$ its curvature. We consider also the bundle of groups $\mathrm{Int}P = P \times_G G$ (G acts on itself by internal automorphisms: $h \mapsto ghg^{-1}, \ h, g \in G$) associated with P, the bundle of Lie algebras $\mathrm{Ad}P = P \times_G \mathfrak{g}$ and a complex vector bundle $E = P \times_G V$, where V is the space of some irreducible representation of G. All these associated bundles inherit their connection $\mathcal{A}$ from P. For the simplicity one can consider $G = \mathrm{SU}(k)$, $\mathfrak{g} = \mathrm{su}(k)$ and $V = \mathbb{C}^k$

We denote by $\mathbb{A}'$ the space of connections on P and by $\mathcal{G}'$ the infinite-dimensional group of gauge transformations (automorphisms of P which induce the identity transformation of X),

$$\mathcal{A} \ \mapsto \ \mathcal{A}^g \ = \ g^{-1} \mathcal{A} g + g^{-1} \mathrm{d} g \ , \tag{2.1}$$

which can be identified with the space $\Gamma(X, \mathrm{Int}P)$ of global sections of the bundle $\mathrm{Int}P$. Correspondingly, the infinitesimal action of $\mathcal{G}'$ is defined by global sections χ of the bundle $\mathrm{Ad}P$,

$$\mathcal{A} \ \mapsto \ \delta_\chi \mathcal{A} \ = \ \mathrm{d}\chi + [\mathcal{A}, \chi] \ =: \ D_{\mathcal{A}} \chi \tag{2.2}$$

with $\chi \in \mathrm{Lie}\, \mathcal{G}' = \Gamma(X, \mathrm{Ad}P)$.

We restrict ourselves to the subspace $\mathbb{A} \subset \mathbb{A}'$ of irreducible connections and to the subgroup $\mathcal{G} = \mathcal{G}' / Z(\mathcal{G}')$ of $\mathcal{G}'$ which acts freely on $\mathbb{A}$. Then the *moduli space* of irreducible connections on P

[1] In this section we follow the discussion of [21].

(and on E) is defined as the quotient $\mathbb{A}/\mathcal{G}$. Classes of gauge equivalent connections are points $[\mathcal{A}]$ in $\mathbb{A}/\mathcal{G}$.

Since $\mathbb{A}$ is an affine space, for each $\mathcal{A} \in \mathbb{A}$ we have a canonical identification between the tangent space $T_{\mathcal{A}}\mathbb{A}$ and the space $\Lambda^1(X, \mathrm{Ad}P)$ of 1-forms on X with values in the vector bundle $\mathrm{Ad}P$. Our $\mathfrak{g} = \mathrm{su}(k)$ is a matrix Lie algebra, with the metric defined by the trace. The metrics on X and on the Lie algebra $\mathrm{su}(k)$ induce an inner product on $\Lambda^1(X, \mathrm{Ad}P)$,

$$\langle \xi_1, \xi_2 \rangle \;=\; \int_X \mathrm{tr}\,(\xi_1 \wedge *\xi_2) \qquad \text{for} \qquad \xi_1, \xi_2 \in \Lambda^1(X, \mathrm{Ad}P)\ . \tag{2.3}$$

This inner product is transferred to $T_{\mathcal{A}}\mathbb{A}$ by the canonical identification. It is invariant under the $\mathcal{G}$-action on $\mathbb{A}$, whence we get a metric (2.3) on the moduli space $\mathbb{A}/\mathcal{G}$.

Instanton connections. Suppose there exists a $(d-4)$-form Σ on X which allows us to introduce the instanton equation

$$*\mathcal{F}_X + \Sigma \wedge \mathcal{F}_X \;=\; 0 \tag{2.4}$$

discussed in Section 1. We denote by $\mathcal{N} \subset \mathbb{A}$ the space of irreducible connections subject to (2.4) on the rank-k complex vector bundle $E \to X$. This space $\mathcal{N}$ of instanton solutions on X is a subspace of the affine space $\mathbb{A}$, and we define the moduli space $\mathcal{M}$ of instantons as the quotient space

$$\mathcal{M} \;=\; \mathcal{N}/\mathcal{G} \tag{2.5}$$

together with a projection

$$\pi :\ \mathcal{N} \stackrel{\mathcal{G}}{\to} \mathcal{M}\ . \tag{2.6}$$

According to the bundle structure (2.6), at any point $\mathcal{A} \in \mathcal{N}$, the tangent bundle $T_{\mathcal{A}}\mathcal{N} \to \mathcal{N}$ splits into the direct sum

$$T_{\mathcal{A}}\mathcal{N} \;=\; \pi^* T_{[\mathcal{A}]}\mathcal{M} \oplus T_{\mathcal{A}}\mathcal{G}\ . \tag{2.7}$$

In other words,

$$T_{\mathcal{A}}\mathcal{N} \ni\ \tilde{\xi} \;=\; \xi + D_{\mathcal{A}}\chi \qquad \text{with} \qquad \xi \in \pi^* T_{[\mathcal{A}]}\mathcal{M} \quad \text{and} \quad D_{\mathcal{A}}\chi \in T_{\mathcal{A}}\mathcal{G}\ , \tag{2.8}$$

where $\tilde{\xi}, \xi \in \Lambda^1(X, \mathrm{Ad}P)$ and $\chi \in \Lambda^0(X, \mathrm{Ad}P) = \Gamma(X, \mathrm{Ad}P)$. The choice of ξ corresponds to a local fixing of a gauge. We denote by ξ_α a local basis of vector fields on $\mathcal{M}$ (sections of the tangent bundle $T\mathcal{M}$) with $\alpha = 1, \ldots, \dim_{\mathbb{R}}\mathcal{M}$. Restricting the metric (2.3) on $\mathbb{A}/\mathcal{G}$ to the subspace $\mathcal{M}$ provides a metric $\mathbb{G} = (G_{\alpha\beta})$ on the instanton moduli space,

$$G_{\alpha\beta} \;=\; \int_X \mathrm{tr}\,(\xi_\alpha \wedge *\xi_\beta)\ . \tag{2.9}$$

Using this metric on $\mathcal{M}$, we can introduce Christoffel symbols

$$\Gamma^\gamma_{\alpha\beta} = \frac{1}{2}\, G^{\gamma\varkappa}\big(\partial_\alpha G_{\beta\varkappa} + \partial_\beta G_{\alpha\varkappa} - \partial_\varkappa G_{\alpha\beta}\big)\ , \tag{2.10}$$

where the derivatives $\partial_\alpha := \partial/\partial\phi^\alpha$ are taken with respect to local coordinates ϕ^α on $\mathcal{M}$ in which $\mathbb{G} = G_{\alpha\beta}\,\mathrm{d}\phi^\alpha\,\mathrm{d}\phi^\beta$. One can also introduce Riemannian tensor, Ricci tensor etc.

3 Adiabatic limit for the Yang-Mills equations in $d \geq 4$

Splitting of the Yang-Mills equations. So, we consider the manifold

$$M = \mathbb{R} \times X \tag{3.1}$$

with a metric

$$\hat{g}_\varepsilon = -\mathrm{d}t^2 + \varepsilon^2 g = -\mathrm{d}t^2 + \varepsilon^2 g_{\mu\nu}\mathrm{d}x^\mu \mathrm{d}x^\nu \ , \tag{3.2}$$

and rank-k complex vector bundle $E \to M$ with an $\mathrm{su}(k)$-valued connection $\mathcal{A}$ as well as the curvature 2-form

$$\hat{\mathcal{F}} = \tfrac{1}{2}\,\mathcal{F}_{\hat{\mu}\hat{\nu}}\mathrm{d}x^{\hat{\mu}} \wedge \mathrm{d}x^{\hat{\nu}} = \mathcal{F}_{0\mu}\mathrm{d}x^0 \wedge \mathrm{d}x^\mu + \tfrac{1}{2}\,\mathcal{F}_{\mu\nu}\mathrm{d}x^\mu \wedge \mathrm{d}x^\nu \ . \tag{3.3}$$

Recall that we assume that the second part in (3.3),

$$\mathcal{F}_X = \tfrac{1}{2}\,\mathcal{F}_{\mu\nu}\mathrm{d}x^\mu \wedge \mathrm{d}x^\nu \ , \tag{3.4}$$

satisfies the instanton equation (2.4) and for the connection $\hat{\mathcal{A}}$ on $E \to \mathbb{R} \times X$ we have

$$\hat{\mathcal{A}} = \mathcal{A}_{\hat{\mu}}\mathrm{d}x^{\hat{\mu}} = \mathcal{A}_0\mathrm{d}x^0 + \mathcal{A}_\mu \mathrm{d}x^\mu = \mathcal{A}_0\mathrm{d}t + \mathcal{A} \tag{3.5}$$

where $\mathcal{A}$ has components only along X but depends on all coordinates (t, x^μ) on M.

We assume that $\mathcal{A}$ satisfies the instanton equation (2.4) for any t and depend on t only via moduli ϕ^α (collective coordinates) described in Section 2. On the other hand, the full Yang-Mills equations (1.4) impose restrictions on dynamics of $\phi^\alpha(t)$. In order to find them we note that for the metric (3.2) we have

$$\hat{\mathcal{F}}^{0\mu} = \hat{g}^{00}\hat{g}^{\mu\nu}\mathcal{F}_{0\nu} = \varepsilon^{-2}\mathcal{F}^{0\nu} \ , \quad \hat{\mathcal{F}}^{\mu\nu} = \varepsilon^{-4}\mathcal{F}^{\mu\nu} \ , \tag{3.6}$$

where in $\mathcal{F}^{0\mu}$ and $\mathcal{F}^{\mu\nu}$ indices are raised by g^{00} and $g^{\mu\nu}$. After substitution of (3.6) into (1.4) we obtain the equations

$$D_\mu \mathcal{F}^{\mu 0} \equiv g^{\mu\nu} D_\mu \mathcal{F}_{0\nu} = 0 \ , \tag{3.7}$$

$$g^{\mu\nu} D_0 \mathcal{F}_{0\nu} = 0 \ , \tag{3.8}$$

where we used that $D_\mu \mathcal{F}^{\mu\nu} = 0$ since $\mathcal{A}_\mu$ is an instanton on X.

Projection on $\mathcal{M}$. For $t \in \mathbb{R}$ varying, the connection $\mathcal{A} = \mathcal{A}(\phi^\alpha(t), x^\mu)$ on the bundle $E \to \{t\} \times X$ defines a map

$$\phi : \mathbb{R} \to \mathcal{M} \quad \text{with} \quad \phi(t) = \{\phi^\alpha(t)\} \ , \tag{3.9}$$

where ϕ^α with $\alpha = 1, ..., \dim_{\mathbb{R}}\mathcal{M}$ are local coordinates on $\mathcal{M}$. This map is not free - it is constrained by the equations (3.7)-(3.8). Since $\mathcal{A}$ belongs to the solution space $\mathcal{N}$ of the instanton equation (2.4), its derivative $\partial_0\mathcal{A}$ is a solution of the linearized form of (2.4) around $\mathcal{A}$, i.e. $\partial_0\mathcal{A}$ belongs to the vector space $T_{\mathcal{A}}\mathcal{N}$. Using (2.7), one can decompose $\partial_0\mathcal{A}_\mu$ into two parts,

$$T_{\mathcal{A}}\mathcal{N} = \pi^* T_{[\mathcal{A}]}\mathcal{M} \oplus T_{\mathcal{A}}\mathcal{G} \quad \Leftrightarrow \quad \partial_0\mathcal{A}_\mu = (\partial_0\phi^\alpha)\xi_{\alpha\mu} + D_\mu\epsilon_0 \ , \tag{3.10}$$

where $\xi_\alpha = \xi_{\alpha\mu} \mathrm{d}x^\mu$ is a local basis of vector fields on $\mathcal{M}$ and ϵ_0 is an su(k)-valued gauge parameter which is determined by the gauge-fixing equations

$$g^{\mu\nu} D_\mu \xi_{\alpha\nu} = 0 \tag{3.11}$$

and therefore from (3.10) and (3.11) we get

$$g^{\mu\nu} D_\mu \partial_0 \mathcal{A}_\nu = g^{\mu\nu} D_\mu D_\nu \epsilon_0 \ . \tag{3.12}$$

Note that

$$\mathcal{F}_{\nu 0} = D_\nu \mathcal{A}_0 - D_0 \mathcal{A}_\nu \ . \tag{3.13}$$

Let us fix the gauge of the Yang-Mills fields on $\mathbb{R} \times X$ by choosing

$$\mathcal{A}_0 := \epsilon_0 \ . \tag{3.14}$$

Then from (3.10) we obtain

$$\mathcal{F}_{\nu 0} = -\dot{\phi}^\alpha \xi_{\alpha\nu} \ , \tag{3.15}$$

where we denoted by dot the derivative with respect to time t. From (3.11) and (3.15) we see that the equations (3.7) are satisfied. Furthermore, since

$$\partial_0 \mathcal{A}_\mu = \dot{\phi}^\alpha \frac{\partial \mathcal{A}_\mu}{\partial \phi^\alpha} \ , \tag{3.16}$$

we get from (3.12) that

$$\mathcal{A}_0 = \epsilon_0 = \dot{\phi}^\alpha \epsilon_\alpha \ , \tag{3.17}$$

where the gauge parameters ϵ_α can be obtained as solutions of the equations

$$g^{\mu\nu} D_\mu D_\nu \epsilon_\alpha = g^{\mu\nu} D_\mu \frac{\partial \mathcal{A}_\nu}{\partial \phi^\alpha} \ , \tag{3.18}$$

which follow from (3.12),(3.16) and (3.17). Notice that $\mathcal{F}_{0\mu}$, given in (3.15), is the projection of $\partial_0 \mathcal{A}_\mu$ from $T_{\mathcal{A}}\mathcal{N}$ to $T_{[\mathcal{A}]}\mathcal{M}$ (cf. [8]):

$$\pi_* \partial_0 \mathcal{A}_\mu = \mathcal{F}_{0\mu} = \dot{\phi}^\alpha \xi_{\alpha\mu} \ . \tag{3.19}$$

Geodesics. Although the evolution of the gauge fields does not exactly follow a trajectory $\phi^\alpha(t)$ in the set of exact static solutions (moduli space $\mathcal{M}$ of instantons on X in our case), it does a good approximation. Following [22], we will show that in the adiabatic limit $\varepsilon \to 0$ the approximation becomes exact and $\phi(t)$ is a geodesic motion on $\mathcal{M}$. To show this, we substitute (3.15) in the remaining unsolved equations (3.8) and obtain

$$g^{\mu\nu} \frac{\mathrm{d}}{\mathrm{d}t} \left(\dot{\phi}^\beta \xi_{\beta\nu} \right) = g^{\mu\nu} \dot{\phi}^\beta [\xi_{\beta\nu}, \epsilon_0] \ . \tag{3.20}$$

Now let us multiply these equations on $\dot{\phi}^\alpha \xi_{\alpha\mu}$, take trace tr over su($k$) and integrate over X. We get the equations[2]

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(G_{\alpha\beta}\, \dot{\phi}^\alpha\, \dot{\phi}^\beta \right) = 0 \tag{3.21}$$

[2]The right hand side of (3.20) vanishes since $g^{\mu\nu} \dot{\phi}^\alpha \dot{\phi}^\beta \mathrm{tr}([\xi_{\alpha\mu}, \xi_{\beta\nu}], \epsilon_0) \equiv 0$ due to converting symmetric and antisymmetric in $(\alpha\beta)$ parts.

on the moduli space $\mathcal{M}$. In deriving (3.21) we identify t with the affine parameter s entering in definition of the metric

$$\mathrm{d}s^2 = G_{\alpha\beta}\,\mathrm{d}\phi^\alpha\,\mathrm{d}\phi^\beta \tag{3.22}$$

on $\mathcal{M}$, where the metric components $G_{\alpha\beta}$ were introduced in (2.3):

$$G_{\alpha\beta} = \int g^{\mu\nu}\,\mathrm{tr}(\xi_{\alpha\mu}\wedge *\xi_{\beta\nu}) \tag{3.23}$$

with the Hodge operator $*$ on X.

Equation (3.21) defines geodesics on $\mathcal{M}$. To see them in more standard form, with Christoffel symbols

$$\Gamma^\alpha_{\beta\gamma} = G^{\alpha\lambda}\left(\frac{\partial}{\partial\phi^\gamma}\,G_{\beta\lambda} + \frac{\partial}{\partial\phi^\beta}\,G_{\alpha\lambda} - \frac{\partial}{\partial\phi^\lambda}\,G_{\alpha\beta}\right)\,, \tag{3.24}$$

we consider the action functional

$$\tilde{S} = \int \mathrm{d}t\sqrt{G_{\alpha\beta}\dot{\phi}^\alpha\,\dot{\phi}^\beta}\,. \tag{3.25}$$

The Euler-Lagrange equations for (3.25) are

$$\ddot{\phi}^\alpha + \Gamma^\alpha_{\beta\gamma}\dot{\phi}^\beta\dot{\phi}^\gamma - \dot{\phi}^\alpha\frac{\mathrm{d}}{\mathrm{d}t}\ln(G_{\beta\gamma}\dot{\phi}^\beta\dot{\phi}^\gamma) = 0 \quad \overset{(3.21)}{\Longrightarrow} \quad \ddot{\phi}^\alpha + \Gamma^\alpha_{\beta\gamma}\,\dot{\phi}^\beta\,\dot{\phi}^\gamma = 0 \tag{3.26}$$

In other words, (3.20)-(3.21) yield equations (3.26) of geodesics on the moduli space $\mathcal{M}$ of instantons on X. This also reflects the well-known (classical) equivalence of the action functional (3.25) and the functional

$$S = \int \mathrm{d}t\,G_{\alpha\beta}\dot{\phi}^\alpha\,\dot{\phi}^\beta \tag{3.27}$$

for which (3.26) are the Euler-Lagrange equations. Note that (3.27) is the effective action for the standard Yang-Mills action functional on $\mathbb{R}\times X$ in the limit $\varepsilon\to 0$. It stems from the term

$$\int_M d\,\mathrm{vol}\,\mathrm{tr}(\mathcal{F}_{0\mu}\mathcal{F}^{0\mu}) \tag{3.28}$$

Finally, note that the pair $(\mathcal{A}_0(\phi(t)), \mathcal{A}_\mu(\phi(t),x))$ can be understood as a connection on $\mathbb{R}\times X$ which obeys part of the Yang-Mills equations and in the neighbourhood of $(\mathcal{A}_0,\mathcal{A}_\mu)$ there is a solution of the full Yang-Mills equations with $\varepsilon\neq 0$ at least for ε sufficiently small (cf. [8, 13, 7]). This follows from the implicit function theorem and means also the bijectivity of moduli space of the time-dependent solutions for $\varepsilon = 0$ and small $\varepsilon\neq 0$ (cf. [13, 7]).

Monopoles and vortices. It is well known that instanton equations on X^d can be reduced to monopole equations on a submanifold X^{d-1} in X^d and similarly (generalized) vortex equations can be obtained by a reduction on a submanifold X^{d-n} with $n\geq 2$. That is why we will be brief and mention only some examples.

Canonical example is given by the case $d=4$. Considering $X^4=\mathbb{R}^4$ and imposing translation invariance with respect to the fourth coordinate x^4 on $\mathbb{R}^4$ one sees that anti-self-dual Yang-Mills equations on $\mathbb{R}^4$ are reduced to the Yang-Mills-Higgs Bogomolny equations on $\mathbb{R}^3$ describing non-abelian monopoles [1, 2, 5, 6]. Then our consideration produces geodesics on the monopole moduli

space reproducing Manton's result [8]. In principle, the same can be done for $d > 4$. For example, monopoles on G_2-holonomy manifolds X^7 can be obtained from Spin(7)-instantons on X^8 as in [23].

Similarly, as was mentioned in the Introduction, the anti-self-dual Yang-Mills equations on the manifold $X^4 = \Sigma_2 \times S^2$ are reduced by imposing SO(3)-symmetry to vortex equations on a Riemannian 2-manifold Σ_2 (see e.g. [2, 3] and references therein). In other words, vortices on Σ_2 can be considered as SO(3)-symmetric instantons on $\Sigma_2 \times S^2$. Then the adiabatic approach to the Yang-Mills equations on $\mathbb{R} \times \Sigma_2 \times S^2$ yields to geodesics on vortex moduli space. The same reduction from instantons to vortices can be done for $d > 4$ (see e.g. [24, 25]) for $X^d = X^{2p} \times X^{2q}$ with Kähler manifolds X^{2p} and X^{2q}. Then one obtains generalized vortex equations (see e.g. [24, 25]) on X^{2p} and the adiabatic approach will describe slowly moving vortices via geodesics on moduli space of vortices on X^{2p} or symmetric instantons on $X^{2p} \times X^{2q}$.

Acknowledgements

This work was partially supported by the Heisenberg-Landau Program.

References

[1] R. Rajaraman, *Solitons and instantons*, North-Holland, Amsterdam, 1984.

[2] A.M. Jaffe and C.H. Taubes, *Vortices and monopoles*, Birkhaeuser, Boston, 1980.

[3] A.D. Popov, "Integrability of vortex equations on Riemann surfaces," Nucl. Phys. B **821** (2009) 452 [arXiv:0712.1756 [hep-th]];

A.D. Popov, "Non-Abelian vortices on Riemann surfaces: an integrable case," Lett. Math. Phys. **84** (2008) 139 [arXiv:0801.0808 [hep-th]].

[4] A.G. Sergeev, *Vortices and Seiberg-Witten equations,* Nagoya Univ. Math. Lectures, Nagoya, 2002.

[5] N.S. Manton and P. Sutcliffe, *Topological solitons,* Cambridge University Press, Cambridge, 2004.

[6] E.J. Weinberg, *Classical solutions in quantum field theory*, Cambridge University Press, Cambridge, 2012.

[7] A.G. Sergeev, "Adiabatic limit in the Ginzburg-Landau and Seiberg-Witten equations", Proc. Steklov Math. Inst. **289** (2015) 227.

[8] N.S. Manton, "A remark on the scattering of BPS monopoles," Phys. Lett. B **110** (1982) 54.

[9] A.G. Sergeev and S.V. Chechin, "Scattering of slowly moving vortices in the Abelian (2+1)-dimensional Higgs model," Theor. Math. Phys. **85** (1990) 1289.

[10] M. Eto, T. Fujimori, T. Nagashima, M. Nitta, K. Ohashi and N. Sakai, "Dynamics of domain wall networks," Phys. Rev. D **76** (2007) 125025 [arXiv:0707.3267 [hep-th]].

[11] K. Peeters and M. Zamaklar, "Motion on moduli spaces with potentials," JHEP **12** (2001) 032 [hep-th/0107164];

J.P. Allen and D.J. Smith, "The low energy dynamics of charge two dyonic instantons," JHEP **02** (2013) 113 [arXiv:1210.3208 [hep-th]].

[12] M. Eto, T. Fujimori, M. Nitta, K. Ohashi and N. Sakai, "Higher derivative corrections to non-Abelian vortex effective theory," Prog. Theor. Phys. **128** (2012) 67 [arXiv:1204.0773 [hep-th]].

[13] D. Stuart, "Dynamics of Abelian Higgs vortices in the near-Bogomolny regime," Commun. Math. Phys. **159** (1994) 51;

D. Stuart, "The geodesic approximation for the Yang-Mills-Higgs equations," Commun. Math. Phys. **166** (1994) 149.

[14] E. Corrigan, C. Devchand, D.B. Fairlie and J. Nuyts, "First order equations for gauge fields in spaces of dimension greater than four," Nucl. Phys. B **214** (1983) 452.

[15] S.K. Donaldson and R.P. Thomas, "Gauge theory in higher dimensions," in: *The Geometric Universe*, Oxford University Press, Oxford, 1998.

[16] G. Tian, "Gauge theory and calibrated geometry," Ann. Math. **151** (2000) 193 [arXiv:math/0010015 [math.DG]];

[17] D. Harland, T.A. Ivanova, O. Lechtenfeld and A.D. Popov, "Yang-Mills flows on nearly Kähler manifolds and G_2-instantons," Commun.Math.Phys. **300** (2010) 185 [arXiv:0909.2730[hep-th]];

K.-P. Gemmer, O. Lechtenfeld, C. Nölle and A.D. Popov, "Yang-Mills instantons on cones and sine-cones over nearly Kähler manifolds," JHEP **09** (2011) 103 [arXiv:1108.3951 [hep-th]].

[18] D. Harland and A.D. Popov, "Yang-Mills fields in flux compactifications on homogeneous manifolds with SU(4)-structure," JHEP **02** (2012) 107 [arXiv:1005.2837 [hep-th]];

A.D. Popov and R.J. Szabo, "Double quiver gauge theory and nearly Kähler flux compactifications," JHEP **02** (2012) 033 [arXiv:1009.3208 [hep-th]].

[19] D. Harland and C. Nölle, "Instantons and Killing spinors," JHEP **03** (2012) 082 [arXiv:1109.3552 [hep-th]];

T.A. Ivanova and A.D. Popov, "Instantons on special holonomy manifolds," Phys. Rev. D **85** (2012) 105012 [arXiv:1203.2657 [hep-th]].

[20] S. Bunk, T.A. Ivanova, O. Lechtenfeld, A.D. Popov and M. Sperling, "Instantons on sine-cones over Sasakian manifolds," Phys. Rev. D **90** (2014) 065028 [arXiv:1407.2948 [hep-th]];

S. Bunk, O. Lechtenfeld, A.D. Popov and M. Sperling, "Instantons on conical half-flat 6-manifolds," JHEP **01** (2015) 030 [arXiv:1409.0030 [hep-th]].

[21] A. Deser, O. Lechtenfeld and A.D. Popov, "Sigma-model limit of Yang-Mills instantons in higher dimensions," Nucl. Phys. B **894** (2015) 361 [arXiv:1412.4258 [hep-th]].

[22] O. Lechtenfeld and A.D. Popov, "Yang-Mills moduli space in the adiabatic limit," J. Phys. A **48** (2015) 425401 [arXiv:1505.05448 [hep-th]].

[23] G. Oliveira, "Monopoles on the Bryant-Salamon G_2-manifolds," J. Geom. Phys. **86** (2014) 599 [arXiv:1310.7392 [math.DG]];

S.A. Cherkis, "Octonions, monopoles, and knots," Lett. Math. Phys. **105** (2015) 641 [arXiv:1403.6836 [hep-th]].

[24] O. García-Prada, "Invariant connections and vortices," Commun.Math.Phys. **156** (1993) 527;

L. Álvarez-Cónsul and O. García-Prada, "Dimensional reduction, SL(2,$\mathbb{C}$)-equivariant bundles and stable holomorphic chains," Int. J. Math. **12** (2001) 159 [math/0112159 [math-dg]].

L. Álvarez-Cónsul and O. García-Prada, "Dimensional reduction and quiver bundles," J. Reine Angew. Math. **556** (2003) 1 [math/0112160 [math-dg]].

[25] A.D. Popov and R.J. Szabo, "Quiver gauge theory of non-Abelian vortices and noncommutative instantons in higher dimensions," J. Math. Phys. **47** (2006) 012306 [hep-th/0504025];

O. Lechtenfeld, A.D. Popov and R.J. Szabo, "Rank two quiver gauge theory, graded connections and noncommutative vortices," JHEP **09** (2006) 054 [hep-th/0603232].

O. Lechtenfeld, A.D. Popov and R.J. Szabo, "SU(3)-equivariant quiver gauge theories and nonabelian vortices," JHEP **08** (2008) 093 [arXiv:0806.2791 [hep-th]].